



Patro Golab, B., Santos, S., Voerman, E., Lawlor, D., Jaddoe, V. W. V., Gaillard, R. (2018). Influence of maternal obesity on the association between common pregnancy complications and risk of childhood obesity: an individual participant data meta-analysis. *Lancet Child and Adolescent Health*, 2(11), 812-821.
[https://doi.org/10.1016/S2352-4642\(18\)30273-6](https://doi.org/10.1016/S2352-4642(18)30273-6)

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/S2352-4642\(18\)30273-6](https://doi.org/10.1016/S2352-4642(18)30273-6)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

SUPPLEMENTAL MATERIALS

Common pregnancy complications and risk of childhood obesity - influence of maternal obesity: An individual participant data meta-analysis

Bernadeta Patro Golab, MD*, Susana Santos, PhD*, Ellis Voerman, MSc, Prof Debbie A. Lawlor, PhD, Prof Vincent W.V. Jaddoe, MD*, Romy Gaillard, MD* on behalf of the Maternal Obesity Childhood Outcomes (MOCO) Study Group †

† Study group members are listed at the end of the Article

* Authors contributed equally

Contents

Figure S1. Flow chart of participating cohorts and individuals.

Table S1. Cohort-specific methods of data collection for pregnancy complications, maternal and childhood anthropometrics, and other covariates.

Table S2. Characteristics of available covariates per cohort.

Table S3. Associations of gestational diabetes, pre-eclampsia and gestational hypertension with offspring BMI in early, mid, and late childhood: models with categorical or continuous maternal BMI and age.

Table S4. Characteristics of the participating pregnancy and birth cohorts.

Table S5. Gestational hypertension, pre-eclampsia, gestational diabetes and the odds ratios of underweight (compared with normal weight) in early, mid, and late childhood.

Table S6. Gestational hypertension, pre-eclampsia, gestational diabetes and the odds ratios of overweight and obesity (compared with normal weight) in early, mid, and late childhood: maternal BMI model and models additionally adjusted for gestational diabetes or gestational hypertension and pre-eclampsia.

Table S7. Associations of gestational diabetes with childhood BMI in early, mid, and late childhood stratified by maternal BMI group.

Table S8. Associations of gestational diabetes, pre-eclampsia and gestational hypertension with offspring BMI in early, mid, and late childhood: models with pre/early-pregnancy BMI or pre-pregnancy BMI.

Figure S2. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in early childhood.

Figure S3. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in mid childhood.

Figure S4. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in late childhood.

Figure S5. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in early childhood.

Figure S6. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in mid childhood.

Figure S7. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in late childhood.

Figure S8. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in early childhood.

Figure S9. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in mid childhood.

Figure S10. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in late childhood.

Acknowledgments (per cohort)

Conflict of interest disclosures

Funding/support (per cohort)

MOCO study group with affiliations

Figure S1. Flow chart of participating cohorts and individuals.




Table S1. Cohort-specific methods of data collection for pregnancy complications, maternal and childhood anthropometrics, and other covariates.^a

Cohort name (country)	Gestational diabetes	Gestational hypertension	Pre-eclampsia	Childhood weight and height	Maternal height	Maternal pre-/early pregnancy weight	Other covariates
ABCD (The Netherlands)	Self-reported or medical records	Self-reported or medical records	Self-reported or medical records	Measured	Self-reported	Self-reported	Self-reported or medical records
ALSPAC (United Kingdom)	Medical records	Medical records	Medical records	Measured	Self-reported	Self-reported	Self-reported/ medical records
AOB/F (Canada)	Medical records	Medical records	Medical records	Reported	Self-reported	Self-reported	Self-reported
BAMSE (Sweden)	Medical Birth Registry	Medical Birth Registry	Medical Birth Registry	Measured	Medical Birth Registry	Medical Birth Registry	Self-reported
BIB (United Kingdom)	OGT test at 27-28 weeks	Medical records	Medical records	Measured	Measured	Measured	Self-reported or medical records
Co.N.ER (Italy)	Self-reported	Self-reported	Self-reported	Reported	Self-reported	Self-reported	Self-reported
DNBC (Denmark)	Self-reported	Self-reported	Medical records	Reported or measured	Self-reported	Self-reported	Self-reported
EDEN (France)	OGT test and medical records	Clinical examination at 24 weeks of gestation and medical records	Medical records	Measured or medical records	Measured	Self-reported	Self-reported
FCOU (Ukraine)	Medical records	Medical records	Medical records	Medical records	Medical records	Medical records	Self-reported or medical records
GASPII (Italy)	Self-reported	Self-reported	Self-reported	Measured	Self-reported	Self-reported	Self-reported
GECKO Drenthe (The Netherlands)	Self-reported	Self-reported	Self-reported	Measured	Self-reported	Self-reported	Self-reported
GENERATION R (The Netherlands)	Medical records or self-reported	Medical records or self-reported	Medical records or self-reported	Measured	Measured	Self-reported	Self-reported
GENERATION XXI (Portugal)	Medical records	Medical records	Medical records	Measured	Measured/ID card	Self-reported	Self-reported
GENESIS (Greece)	Self-reported	Self-reported	NA	Measured	Self-reported	Self-reported	Self-reported
GINIplus (Germany)	Self-reported	NA	NA	Medical records at 4y, measured and reported at 10 and 15y	Self-reported	Self-reported	Self-reported
HUMIS (Norway)	Medical birth registry	Medical birth registry	Medical birth registry	Reported	Self-reported	Self-reported	Medical birth registry or self-reported
INMA (Spain)	Medical records	Medical records	Medical records	Measured	Measured or self-reported	Self-reported	Self-reported
KOALA (The Netherlands)	Self-reported or medical records	Self-reported or medical records	Self-reported or medical records	Reported	Self-reported	Self-reported	Self-reported
Krakow Cohort (Poland)	Self-reported	Self-reported	Self-reported	Measured	Self-reported	Self-reported	Self-reported or medical records
LISApplus (Germany)	Self-reported	NA	NA	Medical records at 4y, measured and reported at 10 and 15y	Self-reported	Self-reported	Self-reported
MoBa (Norway)	Medical records	Medical records	Medical records	Reported	Self-reported	Self-reported	Self-reported
NINFEA (Italy)	Self-reported	Self-reported	Self-reported	Reported	Self-reported	Self-reported	Self-reported

Table S1. Cohort-specific methods of data collection for pregnancy complications, maternal and childhood anthropometrics, and other covariates (continued).^a

Cohort name (country)	Gestational diabetes	Gestational hypertension	Pre-eclampsia	Childhood weight and height	Maternal height	Maternal pre-/early pregnancy weight	Other covariables
PÉLAGIE (France)	Medical records	Medical records	Medical records	Reported	Self-reported	Self-reported	Self-reported
PIAMA (The Netherlands)	Self-reported	Self-reported	Self-reported	Reported and measured (4 and 8y)	Self-reported	Self-reported	Self-reported
Piccolipiù (Italy)	Self-reported	Self-reported	Self-reported	Measured	Self-reported	Self-reported	Self-reported
Project Viva (United States)	Medical records	Medical records	Medical records	Measured	Self-reported	Self-reported	Self-reported
REPRO_PL (Poland)	Medical records	Medical records	Medical records	Measured	Measured	Self-reported	Self-reported or measured
RHEA (Greece)	Self-reported	Measured or self-reported	Self-reported	Medical records or measured	Measured	Self-reported	Self-reported
ROLO (Ireland)	Medical records	NA	NA	Measured	Measured	Measured	Self-reported or measured
SCOPEBASELINE (Ireland)	Measured	Measured	Measured	Measured	Measured	Measured	Self-reported
SEATON (United Kingdom)	NA	Medical records	Medical records	Measured	Measured	Measured	Self-reported or medical records
Slovak PCB study (Slovakia)	Self-reported	Self-reported	Self-reported	Measured	Self-reported	Self-reported	Self-reported
STEPS (Finland)	Medical records	Medical records	Medical records	Measured	Self-reported	Self-reported	Self-reported
SWS (United Kingdom)	Medical records	Medical records	Medical records	Measured	Measured	Measured	Self-reported or medical records

^a Self-reported/reported refers to data obtained either by questionnaire or by interview. NA, not available. OGT, Oral glucose tolerance

Table S2. Characteristics of available covariates per cohort.^a

Cohort name, number of participants, (country)	Maternal Age, (years)		Parity, n (%)		Maternal Education Level, n (%)				Ethnicity, n (%)		Smoking during pregnancy, n (%)		Maternal pre/early-pregnancy BMI, (kg/m ²)		Offspring Sex, n (%)	
	Median (95% range)	Missings	Nulliparous	Missings	Low	Medium	High	Missings	European/White	Missings	Yes	Missings	Median (95% range)	Missings	Male	Missings
ABCD, n=5512, (The Netherlands)	32·0 (20·0, 40·0)	-	3041 (55·2)	-	1147 (20·8)	2029 (36·8)	2295 (41·6)	41 (0·7)	4095 (74·3)	10 (0·2)	605 (11·0)	269 (4·9)	22·2 (17·9, 33·9)	18 (0·3)	2737 (49·7)	-
ALSPAC, n=9041, (United Kingdom)	29·0 (20·0, 38·0)	635 (7·0)	3969 (43·9)	361 (4·0)	5142 (56·9)	2165 (23·9)	1284 (14·2)	450 (5·0)	8386 (92·8)	477 (5·3)	1849 (20·5)	234 (2·6)	22·2 (18·0, 33·6)	635 (7·0)	4583 (50·7)	-
AOB/F, n=1672, (Canada)	31·0 (22·0, 40·0)	34 (2·0)	844 (50·5)	-	132 (7·9)	1262 (75·5)	275 (16·4)	3 (0·2)	1374 (82·2)	2 (0·1)	144 (8·6)	92 (5·5)	23·0 (18·0, 38·2)	19 (1·1)	884 (52·9)	-
BAMSE, n=3329, (Sweden)	30·0 (22·0, 40·0)	-	1861 (55·9)	-	1103 (33·1)	833 (25·0)	1376 (41·3)	17 (0·5)	2904 (87·2)	233 (7·0)	420 (12·6)	-	22·3 (18·2, 31·6)	399 (12)	1688 (50·7)	-
BIB, n=983, (United Kingdom)	27·0 (18·0, 39·0)	1 (0·1)	384 (39·1)	15 (1·5)	241 (24·5)	371 (37·7)	369 (37·5)	2 (0·2)	401 (40·8)	-	144 (14·6)	1 (0·1)	24·7 (17·6, 39·4)	96 (9·8)	462 (47·0)	-
Co.N.ER, n=528, (Italy)	33·9 (25·5, 42·1)	1 (0·2)	235 (44·5)	2 (0·4)	82 (15·5)	240 (45·5)	205 (38·8)	1 (0·2)	524 (99·2)	-	67 (12·7)	-	21·2 (17·7, 30·4)	6 (1·1)	265 (50·2)	-
DNBC, n=40349, (Denmark)	30·3 (22·9, 39·3)	-	19992 (49·5)	26 (0·1)	3107 (7·7)	14694 (36·4)	22428 (55·6)	120 (0·3)	NA	NA	9641 (23·9)	14 (0·0)	22·5 (18·1, 33·6)	712 (1·8)	20749 (51·4)	-
EDEN, n=1139, (France)	29·9 (21·0, 39·7)	-	727 (53·4)	3 (0·2)	313 (23·0)	248 (18·2)	793 (58·3)	7 (0·5)	NA	NA	310 (22·8)	10 (0·7)	22·1 (17·4, 34·8)	31 (2·3)	721 (53·0)	-
FCOU, n=2332, (Ukraine)	23·0 (17·0, 36·0)	-	1495 (64·1)	181 (7·8)	127 (5·4)	1534 (65·8)	528 (22·6)	143 (6·1)	2332 (100·0)	-	178 (7·6)	138 (5·9)	21·8 (17·3, 32·0)	235 (10·1)	1218 (52·2)	-
GASPII, n=570, (Italy)	33·0 (24·0, 41·0)	-	326 (57·2)	-	78 (13·7)	284 (49·8)	208 (36·5)	-	565 (99·1)	1 (0·2)	64 (11·2)	-	21·3 (17·6, 31·1)	2 (0·4)	294 (51·6)	-
GECKO Drenthe, n=2119 (The Netherlands)	31·0 (22·0, 39·0)	1 (0·0)	793 (37·4)	135 (6·4)	1258 (59·4)	750 (35·4)	-	111 (5·2)	1977 (93·3)	96 (4·5)	288 (13·6)	38 (1·8)	23·7 (18·6, 36·7)	157 (7·4)	1062 (50·1)	-
GENESIS, n=2145, (Greece)	30·2 (20·9, 39·0)	197 (9·2)	1200 (56·0)	-	94 (4·4)	989 (46·2)	804 (37·5)	256 (11·9)	NA	NA	350 (16·3)	207 (9·7)	21·8 (17·6, 30·9)	245 (11·4)	1116 (52·1)	-
GENERATION R n=7550, (The Netherlands)	31·0 (19·8, 39·8)	-	4101 (54·3)	149 (2·0)	667 (8·8)	3025 (40·1)	3138 (41·6)	720 (9·5)	4412 (58·4)	289 (3·8)	1600 (21·2)	942 (12·5)	22·8 (18·1, 34·9)	908 (12·0)	3788 (50·2)	-
GENERATION XXI, n=5921, (Portugal)	30·0 (18·0, 40·0)	-	3395 (57·3)	88 (1·5)	1768 (29·9)	2588 (43·7)	1537 (26·0)	28 (0·5)	NA	NA	1285 (21·7)	58 (1·0)	23·0 (18·2, 34·7)	8 (0·1)	3027 (51·1)	-
GINIplus, n=2313, (Germany)	31·0 (24·0, 40·0)	-	NA	NA	276 (11·9)	977 (42·4)	1054 (45·6)	6 (0·3)	NA	NA	257 (11·1)	28 (1·2)	22·0 (18·0, 31·4)	6 (0·3)	1137 (49·2)	-
HUMIS, n=970, (Norway)	30·0 (22·0, 39·0)	-	425 (43·8)	-	96 (9·9)	153 (15·8)	568 (58·6)	153 (15·8)	723 (74·5)	159 (16·4)	97 (10·0)	43 (4·4)	23·3 (18·3, 35·0)	25 (2·6)	488 (50·30)	-
INMA, n=1933, (Spain)	30·0 (22·0, 39·0)	1 (0·1)	1056 (54·6)	2 (0·1)	583 (30·2)	736 (38·1)	594 (30·7)	20 (1·0)	1846 (95·5)	3 (0·2)	341 (17·6)	14 (0·7)	22·5 (18·0, 34·6)	22 (1·1)	992 (51·3)	-
KOALA, n=2061, (The Netherlands)	32·0 (25·0, 40·0)	2 (0·1)	899 (43·6)	43 (2·1)	178 (8·6)	760 (36·9)	1026 (49·8)	97 (4·7)	1993 (96·7)	6 (0·3)	122 (5·9)	6 (0·3)	22·8 (18·5, 33·5)	10 (0·5)	1059 (51·4)	-
Krakow Cohort, n=424, (Poland)	27·5 (20·0, 34·0)	-	270 (63·7)	-	40 (9·4)	161 (38·0)	223 (52·6)	-	424 (100)	-	-	-	21·1 (17·3, 28·0)	2 (0·5)	217 (51·2)	-

Table S2. Characteristics of available covariates per cohort (continued).^a

Cohort name, number of participants, (country)	Maternal Age, (years)		Parity, n (%)		Maternal Education Level, n (%)				Ethnicity, n (%)		Smoking during Pregnancy, n (%)		Maternal pre/early-pregnancy BMI, (kg/m ²)		Offspring Sex, n (%)	
	Median (95% range)	Missings	Nulliparous	Missings	Low	Medium	High	Missings	European/ White	Missings	Yes	Missings	Median (95% range)	Missings	Male	Missings
LISApplus, n=1584, (Germany)	32·0 (23·0, 40·0)	-	687 (43·4)	5 (0·3)	96 (6·2)	565 (35·7)	908 (57·3)	13 (0·8)	NA	NA	202 (12·8)	7 (0·4)	21·7 (17·8, 32·2)	2 (0·1)	829 (52·3)	-
MoBa, n=55008, (Norway)	30·0 (22·0, 39·0)	-	25506 (46·4)	-	14936 (27·2)	24284 (44·1)	14739 (26·8)	1049 (1·9)	NA	NA	3955 (7·2)	5445 (9·9)	23·1 (18·4, 34·7)	98 (0·2)	28200 (51·30)	-
NINFEA, n=1726, (Italy) ^b	33·0 (25·0, 41·0)	-	1172 (67·9)	1 (0·1)	65 (3·8)	575 (33·3)	1067 (61·8)	19 (1·1)	1703 (98·7)	-	137 (7·9)	25 (1·4)	21·4 (17·3, 31·9)	50 (2·9)	880 (51·0)	-
PÉLAGIE, n=738, (France)	30·1 (22·7, 39·5)	-	326 (44·2)	2 (0·3)	106 (14·4)	128 (17·3)	503 (68·2)	1 (0·1)	NA	NA	198 (26·8)	1 (0·1)	21·7 (17·5, 32·4)	6 (0·8)	381 (51·6)	-
PIAMA, n=1815, (The Netherlands)	31·0 (24·0, 38·0)	5 (0·3)	926 (51·0)	-	300 (16·5)	757 (41·7)	754 (41·5)	4 (0·2)	1745 (96·1)	24 (1·3)	243 (13·4)	11 (0·6)	22·2 (18·3, 30·8)	113 (6·2)	918 (50·6)	-
Piccolipiù, n=687, (Italy)	34·0 (24·0, 43·0)	-	430 (62·6)	-	69 (10·0)	263 (38·3)	354 (51·5)	1 (0·1)	678 (98·7)	1 (0·1)	156 (22·7)	-	21·6 (17·6, 31·8)	2 (0·3)	361 (52·5)	-
Project Viva, n=1389, (United States)	32·4 (18·9, 41·2)	-	664 (47·8)	-	437 (31·5)	487 (35·1)	458 (33·0)	7 (0·5)	954 (68·7)	7 (0·5)	144 (10·4)	31 (2·2)	23·5 (18·3, 38·2)	10 (0·7)	711 (51·2)	-
REPRO_PL, n=291, (Poland)	28·0 (20·0, 37·0)	3 (1·0)	167 (57·4)	-	32 (11·0)	102 (35·1)	157 (54·0)	-	291 (100)	-	34 (11·7)	-	21·6 (17·2, 32·8)	8 (2·7)	139 (47·8)	-
RHEA, n=740, (Greece)	30·0 (20·0, 40·0)	3 (0·4)	NA	NA	105 (14·2)	386 (52·2)	242 (32·7)	7 (0·9)	739 (99·9)	1 (0·1)	256 (34·6)	-	23·4 (18·0, 36·4)	10 (1·4)	396 (53·5)	-
ROLO, n=283, (Ireland)	33·3 (24·7, 40·4)	-	-	-	-	50 (17·7)	204 (72·1)	29 (10·2)	280 (98·8)	-	6 (2·1)	-	25·2 (20·0, 38·7)	3 (1·1)	131 (46·3)	-
SCOPEBASELINE, n=1046, (Ireland)	31·0 (22·0, 39·0)	-	1046 (100·0)	-	-	122 (11·7)	921 (88·3)	3 (0·3)	1033 (98·8)	-	229 (21·9)	-	24·0 (19·2, 34·7)	1 (0·1)	531 (50·8)	-
SEATON, n=872, (United Kingdom)	30·4 (19·5, 39·6)	-	324 (37·2)	-	185 (21·2)	251 (28·8)	340 (39·0)	96 (11·0)	NA	NA	317 (36·4)	-	23·9 (18·8, 37·7)	1 (0·1)	439 (50·3)	-
Slovak PCB study, n=524, (Slovakia)	26·0 (19·0, 38·0)	-	208 (39·7)	1 (0·2)	235 (44·8)	254 (48·5)	31 (5·9)	4 (0·8)	436 (83·2)	-	81 (15·5)	20 (3·8)	21·2 (16·7, 31·6)	44 (8·4)	260 (49·6)	-
STEPS, n=297, (Finland)	31·5 (24·2, 40·4)	-	179 (60·3)	-	15 (5·1)	87 (29·3)	193 (65·0)	2 (0·7)	NA	NA	10 (3·4)	2 (0·7)	22·6 (18·3, 34·3)	1 (0·3)	148 (49·8)	-
SWS, n=2646, (United Kingdom)	30·2 (22·7, 36·4)	738 (27·9)	1394 (52·7)	3 (0·1)	292 (11·0)	1578 (59·6)	769 (29·1)	7 (0·3)	2545 (96·2)	1 (0·0)	335 (12·7)	287 (10·8)	24·1 (18·9, 37·4)	25 (0·9)	1367 (51·7)	-

^aNA, not available; “-” for missings refers to data available in all participants (no missings); ^bSubset of participants with 4-years follow-up completed.

^cDistinguishes between those born in Italy and those born outside Italy.

Table S3. Associations of gestational diabetes, pre-eclampsia and gestational hypertension with offspring BMI in early, mid, and late childhood: models with categorical or continuous maternal BMI and age.

	Childhood BMI in standard-deviation scores Difference (95% Confidence Interval)		
	Early childhood 2.0-4.9y	Mid childhood 5.0-9.9y	Late childhood 10.0-17.9y
Model with categorical maternal BMI and age			
Gestational diabetes	0.05 (0.00, 0.10)	0.08 (0.03, 0.13)	-0.09 (-0.22, 0.03)
Pre-eclampsia	-0.12 (-0.16, -0.07)	-0.01 (-0.05, 0.02)	0.00 (-0.14, 0.14)
Gestational hypertension	0.01 (-0.03, 0.06)	0.03 (0.00, 0.06)	0.07 (0.00, 0.13)
Model with continuous maternal BMI and age			
Gestational diabetes	0.03 (-0.02, 0.08)	0.04 (-0.01, 0.09)	-0.12 (-0.24, 0.00)
Pre-eclampsia	-0.14 (-0.18, -0.09)	-0.04 (-0.07, 0.00)	-0.07 (-0.21, 0.06)
Gestational hypertension	-0.01 (-0.05, 0.04)	0.00 (-0.03, 0.03)	0.01 (-0.05, 0.08)

Values are regression coefficients (95% confidence intervals) from multilevel linear mixed effects models that reflect differences in early childhood (2·0 to 4·9 years), mid childhood (5·0 to 9·9 years) and late childhood (10·0 to 17·9 years) BMI SDS for children born to mothers with pregnancy complication, as compared with the reference group (children born to mothers with an uncomplicated pregnancy). Models are adjusted for offspring's sex, maternal age, educational level, ethnicity, parity, smoking during pregnancy and pre-/early-pregnancy BMI.

Table S4. Characteristics of the participating pregnancy and birth cohorts (n=160757).^a

Cohort name, number of participants, (country)	Age (months)			Body mass index (kg/m ²)		
	Early childhood (2·0-4·9y)	Mid childhood (5·0-9·9y)	Late childhood (10·0-17·9y)	Early childhood (2·0-4·9y)	Mid childhood (5·0-9·9y)	Late childhood (10·0-17·9y)
ABCD, n=5512, (The Netherlands)	47·2 (25·5, 54·3)	68·2 (61·6, 82·2)	NA	15·7 (13·5, 19·1)	15·4 (13·2, 19·7)	NA
ALSPAC, n=9041, (United Kingdom)	48·7 (30·8, 49·6)	115·0 (88·0, 119·0)	165·0 (125·0, 171·0)	16·2 (14·0, 19·2)	16·5 (13·7, 23·8)	19·4 (15·1, 29·0)
AOB/F, n=1672, (Canada)	36·0 (35·0, 42·0)	NA	NA	15·7 (12·9, 19·5)	NA	NA
BAMSE, n=3329, (Sweden)	51·4 (48·2, 57·6)	101·0 (89·0, 109·0)	201·3 (191·7, 210·3)	16·1 (14·0, 19·3)	16·7 (14·1, 22·4)	21·2 (17·1, 28·9)
BIB, n=983, (United Kingdom)	36·8 (35·7, 39·2)	NA	NA	16·1 (13·8, 19·3)	NA	NA
Co.N.ER, n=528, (Italy)	43·9 (34·7, 54·7)	95·0 (86·6, 111·1)	NA	15·7 (12·9, 20·0)	16·9 (14·0, 23·0)	NA
DNBC, n=40349, (Denmark)	NA	85·0 (75·5, 89·5)	NA	NA	15·5 (13·0, 19·6)	NA
EDEN, n=1361, (France)	38·0 (36·9, 39·9)	67·7 (65·0, 72·4)	NA	15·8 (13·8, 18·2)	15·2 (13·3, 18·5)	NA
FCOU, n=2332, (Ukraine)	35·0 (24·0, 40·0)	84·0 (75·0, 93·0)	193·0 (183·0, 209·1)	16·3 (13·1, 20·5)	15·4 (12·9, 19·4)	20·4 (16·3, 27·5)
GASPII, n=570, (Italy)	50·0 (43·0, 53·0)	104·0 (98·0, 113·0)	NA	16·3 (14·0, 20·0)	17·1 (13·8, 23·5)	NA
GECKO Drenthe, n=2119, (The Netherlands)	NA	70·4 (62·5, 78·6)	NA	NA	15·8 (13·8, 19·7)	NA
GENESIS, n=2145, (Greece)	43·9 (26·2, 58·1)	62·1 (60·0, 72·1)	NA	16·5 (13·9, 21·4)	16·6 (13·5, 23·2)	NA
GENERATION R, n=7550, (The Netherlands)	45·8 (44·5, 48·6)	115·2 (69·4, 119·4)	122·1 (120·1, 137·8)	15·7 (13·5, 19·2)	16·5 (13·8, 23·8)	17·4 (14·3, 25·5)
GENERATION XXI, n=5921, (Portugal)	50·0 (46·0, 58·0)	85·0 (83·0, 95·0)	NA	16·0 (13·7, 20·6)	16·5 (13·7, 23·5)	NA
GINIplus, n=2313, (Germany)	48·0 (44·0, 52·0)	62·9 (60·1, 75·0)	182·0 (177·0, 191·0)	15·4 (13·2, 18·2)	15·3 (13·1, 18·6)	20·1 (16·0, 27·7)
HUMIS, n=970, (Norway)	25·6 (24·0, 37·3)	84·0 (60·0, 92·0)	NA	16·1 (13·5, 19·2)	15·5 (12·8, 19·7)	NA
INMA, n=1933, (Spain)	52·9 (49·0, 56·5)	83·7 (75·2, 94·5)	174·5 (171·9, 181·5)	16·0 (13·7, 20·0)	16·4 (13·6, 23·5)	20·6 (16·4, 30·3)
KOALA, n=2061, (The Netherlands)	55·5 (48·1, 59·8)	106·0 (61·5, 119·3)	121·4 (120·0, 126·7)	15·1 (12·8, 18·1)	15·6 (13·0, 20·2)	16·2 (13·1, 22·7)
Krakow Cohort, n=424, (Poland)	48·0 (36·0, 51·2)	108·0 (60·0, 111·0)	NA	15·3 (12·9, 18·7)	16·2 (13·5, 22·4)	NA
LISApplus, n=1584, (Germany)	48·0 (44·0, 52·0)	62·7 (60·1, 73·5)	181·0 (173·0, 190·0)	15·4 (13·2, 18·2)	15·1 (13·1, 18·1)	19·9 (16·0, 28·0)
MoBa, n=55008, (Norway)	36·4 (25·4, 59·9)	86·9 (60·9, 100·9)	121·9 (120·9, 122·9)	15·9 (13·2, 19·2)	15·7 (12·9, 20·6)	NA
NINFEA, n=1726, (Italy)	49·7 (48·3, 57·0)	86·1 (84·8, 93·0)	NA	15·4 (12·6, 19·5)	15·4 (12·8, 20·8)	NA
PÉLAGIE, n=738, (France)	24·4 (24·0, 26·4)	NA	NA	16·0 (13·9, 18·7)	NA	NA
PIAMA, n=1815, (The Netherlands)	49·1 (44·3, 54·2)	97·5 (90·8, 109·8)	195·9 (192·4, 203·1)	16·2 (13·9, 19·4)	16·0 (13·5, 21·7)	20·3 (16·8, 27·4)
Piccolipiù, n=687, (Italy)	24·0 (24·0, 28·0)	NA	NA	16·2 (13·4, 19·6)	NA	NA
Project Viva, n=1389, (United States)	37·9 (36·1, 50·2)	92·2 (82·5, 116·5)	123·8 (120·6, 131·1)	16·3 (14·2, 19·5)	16·4 (13·7, 24·4)	18·0 (14·3, 32·9)
REPRO_PL, n=291, (Poland)	25·0 (24·0, 31·0)	88·0 (84·2, 94·0)	NA	16·2 (13·2, 19·6)	16·7 (13·3, 25·2)	NA
RHEA, n=740, (Greece)	49·8 (48·0, 57·8)	NA	NA	16·1 (13·8, 21·3)	NA	NA
ROLO, n=283, (Ireland)	24·7 (24·0, 34·9)	NA	NA	16·0 (13·8, 19·9)	NA	NA
SCOPE BASELINE, n=1046, (Ireland)	25·5 (24·5, 28·9)	NA	NA	16·6 (14·5, 19·3)	NA	NA
SEATON, n=872, (United Kingdom)	58·6 (55·9, 59·9)	61·2 (60·0, 119·7)	180·1 (121·5, 186·1)	16·2 (14·1, 19·6)	16·2 (13·9, 20·0)	20·7 (15·2, 31·0)
Slovak PCB study, n=524, (Slovakia)	45·4 (44·8, 49·7)	93·0 (85·0, 101·0)	NA	18·1 (12·5, 24·9)	16·1 (13·3, 24·1)	NA
STEPS, n=297, (Finland)	36·8 (35·7, 38·4)	NA	NA	16·2 (14·2, 18·8)	NA	NA
SWS, n=2646, (United Kingdom)	38·4 (35·6, 50·7)	80·4 (74·7, 87·1)	NA	16·0 (13·8, 19·3)	15·7 (13·4, 20·8)	NA

^aValues are expressed as medians (95% range). NA, not available.

Table S5. Gestational hypertension, pre-eclampsia, gestational diabetes and the odds ratios of underweight (compared with normal weight) in early, mid, and late childhood.

	Underweight vs normal weight Odds Ratios (95% Confidence Interval)		
	Early childhood 2·0-4·9y	Mid childhood 5·0-9·9y	Late childhood 10·0-17·9y
Basic model			
No gestational hypertension and pre-eclampsia	Reference n _{cases/total} = 1099/64157	Reference n _{cases/total} = 2057/82925	Reference n _{cases/total} = 175/7166
Gestational hypertension	0·83 (0·57, 1·21) n _{cases/total} = 29/2487	0·65 (0·49, 0·86) n _{cases/total} = 50/3650	0·60 (0·35, 1·05) n _{cases/total} = 14/871
Pre-eclampsia	1·33 (1·01, 1·75) n _{cases/total} = 55/2297	1·10 (0·88, 1·38) n _{cases/total} = 80/2806	0·35 (0·09, 1·43) n _{cases/total} = 2/230
No gestational diabetes	Reference n _{cases/total} = 1255/76340	Reference n _{cases/total} = 2237/94444	Reference n _{cases/total} = 287/13029
Gestational diabetes	0·76 (0·48, 1·21) n _{cases/total} = 19/1681	0·80 (0·49, 1·32) n _{cases/total} = 16/1192	0·72 (0·23, 2·26) n _{cases/total} = 3/197

Values are odds ratios (95% confidence intervals) from multilevel binary logistic mixed effects models that reflect the odds of underweight in early, mid and late childhood for offspring exposed to gestational diabetes, gestational hypertension or pre-eclampsia, compared with the reference group (no gestational diabetes/ no gestational hypertension and pre-eclampsia). Results are unadjusted (Basic model). We were unable to further explore adjustment for maternal BMI because of small numbers.

Table S6. Gestational hypertension, pre-eclampsia, gestational diabetes and the odds ratios of overweight and obesity (compared with normal weight) in early, mid, and late childhood: maternal BMI model and models additionally adjusted for gestational diabetes or gestational hypertension and pre-eclampsia.

	Overweight and obesity vs normal weight Odds Ratios (95% Confidence Interval)		
	Early childhood 2·0-4·9y	Mid childhood 5·0-9·9y	Late childhood 10·0-17·9y
Maternal BMI model			
Gestational hypertension	1·01 (0·86, 1·19)	1·02 (0·95, 1·10)	1·18 (1·03, 1·36)
Pre-eclampsia	0·87 (0·73, 1·03)	1·04 (0·95, 1·13)	0·98 (0·71, 1·33)
Maternal BMI model additionally adjusted for gestational diabetes			
Gestational hypertension	1·00 (0·86, 1·18)	1·02 (0·94, 1·09)	1·18 (1·03, 1·36)
Pre-eclampsia	0·87 (0·73, 1·03)	1·04 (0·95, 1·13)	0·98 (0·72, 1·34)
Maternal BMI model additionally adjusted for gestational hypertension and pre-eclampsia			
Gestational diabetes	1·35 (1·15, 1·58)	1·12 (1·00, 1·25)	0·96 (0·71, 1·31)
Gestational diabetes	1·35 (1·15, 1·58)	1·12 (1·00, 1·25)	0·96 (0·71, 1·31)

Values are odds ratios (95% confidence intervals) from multilevel binary logistic mixed effects models that reflect the odds of overweight and obesity in early, mid and late childhood for offspring exposed to gestational diabetes, gestational hypertension or pre-eclampsia, compared with the reference group (no gestational diabetes/ no gestational hypertension and pre-eclampsia). The models are adjusted for offspring's sex, maternal age, educational level, ethnicity, parity, smoking during pregnancy, maternal pre/early-pregnancy BMI (Maternal BMI model), and additionally adjusted for gestational diabetes or gestational hypertension and pre-eclampsia.

Table S7. Associations of gestational diabetes with childhood BMI in early, mid, and late childhood stratified by maternal BMI group.

	Childhood BMI in standard-deviation scores Difference (95% Confidence Interval)		
	Early childhood 2.0-4.9y	Mid childhood 5.0-9.9y	Late childhood 10.0-17.9y
Maternal underweight	0.09 (-0.20, 0.39)	-0.07 (-0.41, 0.27)	-0.59 (-1.30, 0.13)
Maternal normal weight	0.01 (-0.06, 0.08)	0.04 (-0.02, 0.11)	-0.08 (-0.24, 0.09)
Maternal overweight	0.07 (-0.03, 0.17)	0.07 (-0.03, 0.17)	-0.21 (-0.47, 0.05)
Maternal obesity	0.05 (-0.07, 0.16)	0.12 (0.00, 0.24)	0.24 (-0.12, 0.60)


Values are regression coefficients (95% confidence intervals) from multilevel linear mixed effects models that reflect differences in early childhood (2·0 to 4·9 years), mid childhood (5·0 to 9·9 years) and late childhood (10·0 to 17·9 years) BMI SDS for children born to mothers with gestational diabetes, as compared with the reference group (children born to mothers with an uncomplicated pregnancy). Models are adjusted for offspring's sex, maternal age, educational level, ethnicity, parity, and smoking during pregnancy.

Table S8. Associations of gestational diabetes, pre-eclampsia and gestational hypertension with offspring BMI in early, mid, and late childhood: models with pre/early-pregnancy BMI or pre-pregnancy BMI.

	Childhood BMI in standard-deviation scores Difference (95% Confidence Interval)		
	Early childhood 2.0-4.9y	Mid childhood 5.0-9.9y	Late childhood 10.0-17.9y
Model with pre/early-pregnancy BMI			
Gestational diabetes	0.04 (0.00, 0.09)	0.09 (0.04, 0.13)	-0.07 (-0.19, 0.06)
Pre-eclampsia	-0.11 (-0.16, -0.07)	0.00 (-0.04, 0.03)	0.01 (-0.12, 0.14)
Gestational hypertension	0.01 (-0.03, 0.06)	0.04 (0.01, 0.07)	0.07 (0.01, 0.13)
Model with pre-pregnancy BMI			
Gestational diabetes	0.06 (0.01, 0.11)	0.08 (0.03, 0.13)	-0.06 (-0.20, 0.08)
Pre-eclampsia	-0.12 (-0.16, -0.07)	-0.01 (-0.04, 0.03)	0.01 (-0.14, 0.15)
Gestational hypertension	0.02 (-0.02, 0.06)	0.04 (0.01, 0.07)	0.08 (0.01, 0.15)


Values are regression coefficients (95% confidence intervals) from multilevel linear mixed effects models that reflect differences in early childhood (2·0 to 4·9 years), mid childhood (5·0 to 9·9 years) and late childhood (10·0 to 17·9 years) BMI SDS for children born to mothers with pregnancy complication, as compared with the reference group (children born to mothers with an uncomplicated pregnancy). Models are adjusted for offspring's sex, maternal age, educational level, ethnicity, parity, smoking during pregnancy and BMI.

Figure S2. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in early childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in early childhood (2.0-4.9 years) BMI SDS in offspring exposed to gestational diabetes as compared to reference group (no gestational diabetes), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S3. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in mid childhood.




^aValues are pooled differences (95% Confidence Interval, CI) in mid childhood (5.0-9.9 years) BMI SDS in offspring exposed to gestational diabetes as compared to reference group (no gestational diabetes), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S4. 2-stage meta-analysis – association of gestational diabetes with BMI SDS in late childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in late childhood (10.0-17.9 years) BMI SDS in offspring exposed to gestational diabetes as compared to reference group (no gestational diabetes), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S5. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in early childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in early childhood (2.0-4.9 years) BMI SDS in offspring exposed to gestational hypertension as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S6. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in mid childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in mid childhood (5.0-9.9 years) BMI SDS in offspring exposed to gestational hypertension as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S7. 2-stage meta-analysis – association of gestational hypertension with BMI SDS in late childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in late childhood (10.0-17.9 years) BMI SDS in offspring exposed to gestational hypertension as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S8. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in early childhood.^a




^aValues are pooled differences (95% Confidence Interval, CI) in early childhood (2.0-4.9 years) BMI SDS in offspring exposed to pre-eclampsia as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S9. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in mid childhood.^a



^aValues are pooled differences (95% Confidence Interval, CI) in mid childhood (5.0-9.9 years) BMI SDS in offspring exposed to pre-eclampsia as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

Figure S10. 2-stage meta-analysis – association of pre-eclampsia with BMI SDS in late childhood.^a



^aValues are pooled differences (95% Confidence Interval, CI) in late childhood (10.0-17.9 years) BMI SDS in offspring exposed to pre-eclampsia as compared to reference group (no gestational hypertension and pre-eclampsia), expressed by regression coefficients. Analyses are based on maternal BMI model.

ACKNOWLEDGMENTS (PER COHORT)

ABCD

The authors especially thank all participating mothers and their children, and are grateful to all obstetric care providers in Amsterdam for their contribution to the data collection of the ABCD-study.

ALSPAC

The authors are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

AOB/F

The authors acknowledge the contribution and support of All Our Families participants and team.

BAMSE

The authors thank all the children and their parents for participating in the BAMSE cohort and the nurses and other staff members working in the BAMSE project.

BiB

The authors acknowledge that the BiB study is only possible because of the enthusiasm and commitment of the participating children and parents. The authors are grateful to all participants, health professionals and researchers who have made BiB happen.

DNBC

The authors thank all the families for participating in the Danish National Birth Cohort.

EDEN

The authors thank the EDEN mother-child cohort study group, whose members are: I. Annesi-Maesano, J.Y. Bernard, J. Botton, M.A. Charles, P. Dargent-Molina, B. de Lauzon-Guillain, P. Ducimetière, M. de Agostini, B. Foliguet, A. Forhan, X. Fritel, A. Germa, V. Goua, R. Hankard, B. Heude, M. Kaminski, B. Larroque†, N. Lelong, J. Lepeule, G. Magnin, L. Marchand, C. Nabet, F. Pierre, R. Slama, M.J. Saurel-Cubizolles, M. Schweitzer, O. Thiebaugeorges.

FCOU

The authors wish to acknowledge the University of Illinois at Chicago School of Public Health's Louise Hamilton Kyiv Data Management Center for their assistance in the data management for FCOU study.

GASPII

The authors acknowledge the families involved in the study.

GECKO Drenthe

The authors are grateful to the families who took part in the GECKO Drenthe study, the midwives, gynecologists, nurses and GPs for their help for recruitment and measurement of participants, and the whole team from the GECKO Drenthe study.

Generation R

The authors gratefully acknowledge the contribution of general practitioners, hospitals, midwives, and pharmacies in Rotterdam.

Generation XXI

The authors gratefully acknowledge the families enrolled in Generation XXI for their kindness, all members of the research team for their enthusiasm and perseverance and the participating hospitals and their staff for their help and support.

GENESIS

The authors thank the Genesis research group which was comprised from Evdokia Oikonomou, Vivian Detopoulou, Christine Kortsalioudaki, Margarita Bartsota, Thodoris Liargkovinos and Christos Vassilopoulos.

GINIplus

The authors thank all the families for their participation in the GINIplus study. Furthermore, the authors thank all members of the GINIplus Study Group for their excellent work. The GINIplus Study group consists of the following: Institute of Epidemiology I, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg (Heinrich J, Brüske I, Schulz H, Flexeder C, Zeller C, Standl M, Schnappinger M, Ferland M, Thiering E, Tiesler C); Department of Pediatrics, Marien-Hospital, Wesel (Berdel D, von Berg A); Ludwig-Maximilians-University of Munich, Dr von Hauner Children's Hospital (Koletzko S); Child and Adolescent Medicine, University Hospital rechts der Isar of the Technical University Munich (Bauer CP, Hoffmann U); IUF-Environmental Health Research Institute, Düsseldorf (Schikowski T, Link E, Klümper C, Krämer U, Sugiri D).

HUMIS

The authors thank the mothers who participated in the study and the Norwegian Research Council for their continuous support through several grants.

INMA-Valencia

The authors would particularly like to thank all the participants for their generous collaboration.

INMA-Gipuzkoa

The authors thank the children and parents who participated to the INMA-Gipuzkoa study.

INMA-Menorca

The authors thank all the participants for their generous collaboration. The authors are grateful to Mireia Garcia, Maria Victoria Estraña, Maria Victoria Iturriaga, Cristina Capo and Josep LLuch for their assistance in contacting the families and administering the questionnaires.

KOALA

The authors thank the children and parents who participated to the KOALA study.

Krakow Cohort

The authors acknowledge Jagiellonian University Medical College in Krakow and Columbia University in New York. Principal investigator: Prof. FP Perera; co-investigator: Prof. W Jedrychowski.

LISApplus

The authors thank all the families for their participation in the LISApplus study. Furthermore, the authors thank all members of the LISApplus Study Group for their excellent work. The LISApplus Study group consists of the following: Helmholtz Zentrum München, German Research Center for Environmental Health, Institute of Epidemiology I, Munich (Heinrich J, Schnappinger M, Brüske I, Ferland M, Schulz H, Zeller C, Standl M, Thiering E, Tiesler C, Flexeder C); Department of Pediatrics, Municipal Hospital “St. Georg”, Leipzig (Borte M, Diez U, Dorn C, Braun E); Marien Hospital Wesel, Department of Pediatrics, Wesel (von Berg A, Berdel D, Stiers G, Maas B); Pediatric Practice, Bad Honnef (Schaaf B); Helmholtz Centre of Environmental Research – UFZ, Department of Environmental Immunology/Core Facility Studies, Leipzig (Lehmann I, Bauer M, Röder S, Schilde M, Nowak M, Herberth G, Müller J); Technical University Munich, Department of Pediatrics, Munich (Hoffmann U, Paschke M, Marra S); Clinical Research Group Molecular Dermatology, Department of Dermatology and Allergy, Technische Universität München (TUM), Munich (Ollert M, J. Grosch).

MoBa

The authors are grateful to all the participating families in Norway who take part in this on-going cohort study.

NINFEA

The authors thank all families participating in the NINFEA cohort.

PÉLAGIE

The authors thank the gynecologists, obstetricians, ultrasonographers, midwives, pediatricians, and families who participated in the study.

Piccolipiù

The authors acknowledge the Piccolipiù Working Group and the families involved in the study.

Project Viva

The authors thank the Project Viva mothers, children and families for their ongoing participation.

REPRO_PL

The authors would particularly like to thank all the cohort participants for their collaboration.

RHEA

The authors would particularly like to thank all the cohort participants for their generous collaboration.

ROLO The authors are grateful to the families and healthcare staff who partake in ROLO and ROLO kids.

SCOPE BASELINE

The authors are grateful to the families and healthcare staff who kindly gave their time to the study.

Slovak PCB study

The authors thank the Slovak PCB study participants for their ongoing cooperation.

STEPS

The authors are grateful to all the families who took part in STEPS study.

SWS

The authors are grateful to the women of Southampton who gave their time to take part in the Southampton Women’s Survey and to the research nurses and other staff who collected and processed the data.

CONFLICT OF INTEREST DISCLOSURES

Keith M. Godfrey has received reimbursement for speaking at conferences sponsored by companies selling nutritional products, and is part of an academic consortium that has received research funding from Abbott Nutrition, Nestec and Danone. Debbie A. Lawlor has received support from Roche Diagnostics and Medtronic in relation to biomarker research that is not related to the research presented in this paper. Andrea von Berg has received reimbursement for speaking at symposia sponsored by Nestlé and Mead Johnson, who partly financially supported the 15 yrs follow-up examination of the GINIplus study. The rest of the authors reported no conflicts of interest.

FUNDING/SUPPORT (PER COHORT)

ABCD

This work was supported by the Netherlands Organization for Health Research and Development (ZonMw) (TOP grant, 40-00812-98-11010).

ALSPAC

The UK Medical Research Council and Wellcome (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This study has received support from the US National Institute of Health (R01 DK10324) and European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no 669545. DA Lawlor works in a unit that receives UK MRC funding (MC_UU_12013/5) and is an NIHR senior investigator (NF-SI-0611-10196).

AOB/F

All Our Families is funded through Alberta Innovates Interdisciplinary Team Grant #200700595, the Alberta Children's Hospital Foundation, and the Max Bell Foundation.

BAMSE

The Swedish Research Council, The Swedish Heart and Lung Foundation, The Swedish Research Council for Working Life and Social Welfare, the Swedish Asthma and Allergy Association Research Foundation, The Swedish Research Council Formas, Stockholm County Council, and the European Commission's Seventh Framework 29 Program MeDALL under grant agreement No. 261357.

BiB

BiB receives core infrastructure funding from the Wellcome Trust (WT101597MA) and a joint grant from the UK Medical Research Council (MRC) and Economic and Social Science Research Council (ESRC) (MR/N024397/1). This study has received support from the British Heart Foundation (CS/16/4/32482), US National Institute of Health (R01 DK10324) and European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement no 669545. J West is funded by a UK Medical Research Council (MRC) Population Health Scientist Postdoctoral Award (MR/K021656/1). The research was funded by the NIHR CLAHRC Yorkshire and Humber. www.clahrc-yh.nihr.ac.uk and received support from the NIHR Clinical Research Network. The views expressed are those of the author(s), and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Co.N.ER

No funding reported.

DNBC

The Danish National Research Foundation has established the Danish Epidemiology Science Centre that initiated and created the Danish National Birth Cohort. The cohort is furthermore a result of a major grant from this foundation. Additional support for the Danish National Birth Cohort is obtained from the Pharmacy Foundation, the Egmont Foundation, the March of Dimes Birth Defects Foundation, the Augustinus Foundation, and the Health Foundation. The DNBC 7-year follow-up is supported by the Lundbeck Foundation (195/04) and the Danish Medical Research Council (SSVF 0646). The 11-year follow-up was funded by the Danish Independent Research Council, the Lundbeck Foundation and a strategic grant from Department of Public Health, University of Copenhagen.

EDEN

The EDEN study was supported by: Foundation for medical research (FRM), National Agency for Research (ANR), National Institute for Research in Public health (IRES: TGIR cohorte santé 2008 program), French Ministry of Health (DGS), French Ministry of Research, INSERM Bone and Joint Diseases National Research (PRO-A) and

Human Nutrition National Research Programs, Paris-Sud University, Nestlé, French National Institute for Population Health Surveillance (InVS), French National Institute for Health Education (INPES), the European Union FP7 programmes (FP7/2007-2013, HELIX, ESCAPE, ENRIECO, Medall projects), Diabetes National Research Program (through a collaboration with the French Association of Diabetic Patients (AFD)), French Agency for Environmental Health Safety (now ANSES), Mutuelle Générale de l'Éducation Nationale a complementary health insurance (MGEN), French national agency for food security, French speaking association for the study of diabetes and metabolism (ALFEDIAM).

FCOU

FCOU study is supported by the U.S. National Institutes of Health Fogarty International Center, US NIEHS, US CDC, US EPA, and National Academy of Medical Sciences of Ukraine.

GASPII

Ministry of Health.

GECKO Drenthe

The GECKO Drenthe birth cohort was funded by an unrestricted grant of Hutchison Whampoa Ltd, Hong Kong and supported by the University of Groningen, Well Baby Clinic Foundation Icare, Noordlease, Paediatric Association Of The Netherlands and Youth Health Care Drenthe.

Generation R

The general design of the Generation R Study is made possible by financial support from the Erasmus MC, University Medical Center, Rotterdam, Erasmus University Rotterdam, Netherlands Organization for Health Research and Development (ZonMw), Netherlands Organisation for Scientific Research (NWO), Ministry of Health, Welfare and Sport and Ministry of Youth and Families. Research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013), project EarlyNutrition under grant agreement n°289346, the European Union's Horizon 2020 research and innovation programme under grant agreement No 633595 (DynaHEALTH) and the European Union's Horizon 2020 research and innovation programme under grant agreement 733206 (LifeCycle Project). Bernadeta Patro Golab received a research training fellowship grant from the Nestle Nutrition Institute. Romy Gaillard received funding from the Dutch Heart Foundation (grant number 2017T013) and the Dutch Diabetes Foundation (grant number 2017.81.002). Vincent Jaddoe received grants from the Netherlands Organization for Health Research and Development (VIDI 016.136.361) and the European Research Council (Consolidator Grant, ERC-2014-CoG-648916).

Generation XXI

Generation XXI was funded by Programa Operacional de Saúde – Saúde XXI, Quadro Comunitário de Apoio III and Administração Regional de Saúde Norte (Regional Department of Ministry of Health). This study was funded by FEDER through the Operational Programme Competitiveness and Internationalization and national funding from the Foundation for Science and Technology – FCT (Portuguese Ministry of Science, Technology and Higher Education) (POCI-01- 0145-FEDER-016837), under the project “PathMOB.: Risco cardiometabólico na infância: desde o início da vida ao fim da infância” (Ref. FCT PTDC/DTP-EPI/3306/2014) and the Unidade de Investigação em Epidemiologia - Instituto de Saúde Pública da Universidade do Porto (EPIUnit) (POCI-01-0145-FEDER-006862; Ref. UID/DTP/04750/2013). AC Santos holds a FCT Investigator contract IF/01060/2015.

GENESIS

The study was supported by a research grant from Friesland Foods Hellas.

GINIplus

The GINIplus study was mainly supported for the first 3 years of the Federal Ministry for Education, Science, Research and Technology (interventional arm) and Helmholtz Zentrum Munich (former GSF) (observational arm). The 4 year, 6 year, 10 year and 15 year follow-up examinations of the GINIplus study were covered from the respective budgets of the 5 study centres (Helmholtz Zentrum Munich (former GSF), Research Institute at Marien-Hospital Wesel, LMU Munich, TU Munich and from 6 years onwards also from IUF - Leibniz Research-Institute for Environmental Medicine at the University of Düsseldorf) and a grant from the Federal Ministry for Environment (IUF Düsseldorf, FKZ 20462296). Further, the 15 year follow-up examination of the GINIplus study was supported by the Commission of the European Communities, the 7th Framework Program: MeDALL project, and as well by the companies Mead Johnson and Nestlé.

HUMIS

European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreements Early Nutrition n° 289346 and by funds from the Norwegian Research Council's MILPAAHEL programme, project No.213148.

INMA-Sabadell

This study was funded by grants from the Instituto de Salud Carlos III (Red INMA G03/176) and the Generalitat de Catalunya-CIRIT (1999SGR 00241).

INMA-Valencia

This study was funded by Grants from UE (FP7-ENV-2011 cod 282957 and HEALTH.2010.2.4.5-1), Spain: ISCIII (G03/176; FIS-FEDER: PI09/02647, PI11/01007, PI11/02591, PI11/02038, PI13/1944, PI13/2032, PI14/00891, PI14/01687, and PI16/1288; Miguel Servet-FEDER CP11/00178, CP15/00025, and CP116/00051), and Generalitat Valenciana: FISABIO (UGP 15-230, UGP-15-244, and UGP-15-249).

INMA-Gipuzkoa

This study was funded by grants from the Instituto de Salud Carlos III (FISFIS PI06/0867, FIS-PS09/0009) 0867, Red INMA G03/176) and the Departamento de Salud del Gobierno Vasco (2005111093 and 2009111069) and the Provincial Government of Guipúzcoa (DFG06/004 and FG08/001).

INMA-Menorca

This study was funded by grants from the Instituto de Salud Carlos III (Red INMA G03/176).

KOALA

Data collection for the KOALA study from pregnancy up to age 1y was financially supported by grants from Royal Friesland Foods (Leeuwarden); Triodos Foundation (Zeist); Phoenix Foundation; Raphaël Foundation; Iona Foundation; Foundation for the Advancement of Heilpädagogie; Netherlands Organisation for Health Research and Development (ZonMw no. 2100.0090); Netherlands Asthma Foundation (grants no. 3.2.03.48 and 3.2.07.022); Netherlands Heart Foundation (grant no. 2008B112). The funders had no influence on the design, analysis and reporting of the current study.

Krakow Cohort

The study received funding from a NIEHS R01 grants entitled: “Vulnerability of the Fetus/Infant to PAH, PM2.5 and ETS” and “Developmental effects of early-life exposure to airborne PAH” (R01ES010165 and R01ES015282) and from The Lundin Foundation, The John and Wendy Neu Family Foundation, The Gladys and Roland Harriman Foundation and an Anonymous Foundation.

LISApplus

The LISApplus study was mainly supported by grants from the Federal Ministry for Education, Science, Research and Technology and in addition from Helmholtz Zentrum Munich (former GSF), Helmholtz Centre for Environmental Research - UFZ, Leipzig, Research Institute at Marien-Hospital Wesel, Pediatric Practice, Bad Honnef for the first 2 years. The 4 year, 6 year, 10 year and 15 year follow-up examinations of the LISApplus study were covered from the respective budgets of the involved partners (Helmholtz Zentrum Munich (former GSF), Helmholtz Centre for Environmental Research - UFZ, Leipzig, Research Institute at Marien-Hospital Wesel, Pediatric Practice, Bad Honnef, IUF – Leibniz-Research Institute for Environmental Medicine at the University of Düsseldorf) and in addition by a grant from the Federal Ministry for Environment (IUF Düsseldorf, FKZ 20462296). Further, the 15 year follow-up examination of the LISApplus study was supported by the Commission of the European Communities, the 7th Framework Program: MeDALL project.

MoBa

The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research, NIH/NIEHS (contract no N01-ES-75558), NIH/NINDS (grant no.1 U01 NS 047537-01 and grant no.2 U01 NS 047537-06A1).

NINFEA

The NINFEA cohort was partially funded by the Compagnia San Paolo Foundation and by the Piedmont Region.

PÉLAGIE

The Pélagie cohort was supported by the French National Research Agency (ANR-2010-PRSP-007), and the French Research Institute for Public Health (AMC11004NSA-DGS).

PIAMA

The PIAMA study was supported by the Netherlands Organization for Health Research and Development; The Netherlands Organization for Scientific Research; The Netherlands Asthma Fund; The Netherlands Ministry of Spatial Planning, Housing, and the Environment; and The Netherlands Ministry of Health, Welfare, and Sport.

Piccolipiù

The Piccolipiù project was financially supported by the Italian National Center for Disease Prevention and Control (CCM grants years 2010 and 2014) and by the Italian Ministry of Health (art 12 and 12 bis D.lgs 502/92).

Project Viva

National Institutes of Health (R01 HD034568, UG3OD023286).

REPRO_PL

National Science Centre, Poland, under the grant DEC-2014/15/B/NZ7/00998, FP7 HEALS Grant N° 603946 and the Ministry of Science and Higher Education under grant agreement no. 3068/7.PR/2014/2.

RHEA

The "Rhea" project was financially supported by European projects (EU FP6-2003-Food-3-NewGeneris, EU FP6. STREP Hiwate, EU FP7 ENV.2007.1.2.2.2. Project No 211250 Escape, EU FP7-2008-ENV-1.2.1.4 Envirogenomarkers, EU FP7-HEALTH-2009- single stage CHICOS, EU FP7 ENV.2008.1.2.1.6. Proposal No 226285 ENRIECO, EU- FP7- HEALTH-2012 Proposal No 308333 HELIX) and the Greek Ministry of Health (Program of Prevention of obesity and neurodevelopmental disorders in preschool children, in Heraklion district, Crete, Greece: 2011-2014; "Rhea Plus": Primary Prevention Program of Environmental Risk Factors for Reproductive Health, and Child Health: 2012-15).

ROLO

ROLO is supported by the Health Research Board Ireland, the Health Research Centre for Health and Diet Research, and the European Union's Seventh Framework Programme (FP7/2007-2013), project EarlyNutrition under grant agreement no. 289346.

SCOPE BASELINE

The SCOPE Ireland study was funded by the Irish Health Research Board. The Cork BASELINE Birth Cohort Study was funded by the National Children's Research Centre, Dublin, Ireland. SCOPE Ireland was funded by the Health Research Board, Ireland (CSA 2007/2). The BASELINE cohort was supported by the National Children's Research Centre, Dublin, Ireland, and the Food Standards Agency of the United Kingdom (grant no. TO7060). SCOPE and BASELINE are supported by INFANT, an SFI funded Research Centre (grant no 12/RC/2272).

SEATON

The ten year follow up of this cohort was funded by the Medical Research Council (grant 80219), the recruitment and follow up at one, two and five years was funded by Asthma UK (grants 00/011, 02/017).

Slovak PCB study

Support was provided by U.S. National Institutes of Health grants R01 CA096525, R03 TW007152, P30 ES001247, and K12 ES019852.

STEPS

This study was supported by the University of Turku, Abo Akademi University, the Turku University Hospital, and the City of Turku, as well as by the Academy of Finland (grants 121569 and 123571), the Juho Vainio Foundation, the Yrjö Jahnsson Foundation, the Turku.

SWS

The SWS is supported by grants from the Medical Research Council, National Institute for Health Research Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton National Health Service Foundation Trust, and the European Union's Seventh Framework Programme (FP7/2007-2013), project EarlyNutrition (grant 289346). Study participants were drawn from a cohort study funded by the Medical Research Council and the Dunhill Medical Trust.

MOCO Study Group with affiliations

Bernadeta Patro Golab^{*1,2,3}, Susana Santos^{*1,2}, Ellis Voerman^{1,2}, Henrique Barros^{4,5}, Anna Bergström^{6,7}, Marie-Aline Charles^{8,9}, Leda Chatzi^{10,11,12}, Cécile Chevrier¹³, George P. Chrousos¹⁴, Eva Corpeleijn¹⁵, Nathalie Costet¹³, Sarah Crozier¹⁶, Graham Devereux¹⁷, Merete Eggesbø¹⁸, Sandra Ekström⁶, Maria Pia Fantini¹⁹, Sara Farchi²⁰, Francesco Forastiere²⁰, Vagelis Georgiu¹¹, Keith M. Godfrey^{16,21}, Davide Gori¹⁹, Wojciech Hanke²², Irva Hertz-Picciotto²³, Barbara Heude^{8,9}, Daniel Hryhorczuk²⁴, Hazel Inskip^{16,21}, Jesus Ibarluzea^{25,26,27,28}, Louise C. Kenny^{29,30}, Leanne K. Küpers^{15,31,32}, Hanna Lagström³³, Irina Lehmann³⁴, Virissa Lenters¹⁸, Sabrina Llop^{28,35}, Per Magnus³⁶, Renata Majewska³⁷, Johanna Mäkelä³⁸, Yannis Manios³⁹, Fionnuala M. McAuliffe⁴⁰, Sheila W. McDonald⁴¹, John Mehegan⁴², Monique Mommers⁴³, Camilla S. Morgen^{44,45}, George Moschonis⁴⁶, Deirdre Murray^{29,47}, Carol Ni Chaoimh^{29,48}, Ellen A. Nøhr⁴⁹, Anne-Marie Nybo Andersen⁴⁵, Emily Oken⁵⁰, Adriëtte J.J.M. Oostvogels⁵¹, Agnieszka Pac³⁷, Eleni Papadopoulou⁵², Costanza Pizzi⁵³, Kinga Polanska²², Daniela Porta²⁰, Lorenzo Richiardi⁵³, Sheryl L. Rifas-Shiman⁵⁰, Franca Rusconi⁵⁴, Ana C. Santos^{4,5}, Henriette A. Smit⁵⁵, Thorkild I.A. Sørensen^{45,56}, Marie Standl⁵⁷, Camilla Stoltenberg^{58,59}, Jordi Sunyer^{28,60,61}, Michelle Taylor^{31, 32}, Elisabeth Thiering^{57,62}, Carel Thijs⁴³, Maties Torrent⁶³, Suzanne C. Tough^{41,64}, Tomas Trnovec⁶⁵, Steve Turner⁶⁶, Lenie van Rossem⁵⁵, Andrea von Berg⁶⁷, Martine Vrijheid^{28,60,61}, Tanja Vrijkotte⁵¹, Jane West⁶⁸, John Wright⁶⁸, Oleksandr Zvinchuk⁶⁹, Debbie A. Lawlor^{31,32}, Vincent W.V. Jaddoe^{*1,2,70}, Romy Gaillard^{*1,2}

** Authors contributed equally*

¹ The Generation R Study Group, Erasmus MC, University Medical Center, Rotterdam, the Netherlands.

² Department of Pediatrics, Erasmus MC, University Medical Center, Rotterdam, the Netherlands.

³ Department of Pediatrics, Medical University of Warsaw, Warsaw, Poland.

⁴ EPIUnit - Instituto de Saúde Pública, Universidade do Porto, Rua das Taipas, n° 135, 4050-600 Porto, Portugal.

⁵ Department of Public Health and Forensic Sciences and Medical Education, Unit of Clinical Epidemiology, Predictive Medicine and Public Health, University of Porto Medical School, Porto, Portugal.

⁶ Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.

⁷ Centre for Occupational and Environmental Medicine, Stockholm County Council, Stockholm, Sweden.

⁸ INSERM, UMR1153 Epidemiology and Biostatistics Sorbonne Paris Cité Center (CRESS), ORCHAD Team, Villejuif, France.

⁹ Paris Descartes University, Villejuif, France.

¹⁰ Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California, USA.

¹¹ Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Greece.

¹² Department of Genetics and Cell Biology, Maastricht University, the Netherlands.

¹³ Inserm UMR 1085, Irset - Research Institute for Environmental and Occupational Health, F-35000 RENNES, France.

¹⁴ First Department of Pediatrics, Athens University Medical School, Aghia Sophia "Children's Hospital, National and Kapodistrian University of Athens, Athens, Greece.

¹⁵ Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

¹⁶ MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK.

¹⁷ Liverpool School of Tropical Medicine, Liverpool, UK.

¹⁸ Department of Exposure and Environmental Epidemiology, Norwegian Institute of Public Health, Oslo, Norway.

¹⁹ The Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy.

²⁰ Department of Epidemiology, Lazio Regional Health Service, Rome, Italy.

²¹ NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK.

²² Department of Environmental Epidemiology, Nofer Institute of Occupational Medicine, Lodz, Poland.

- ²³ Department of Public Health Sciences, School of Medicine, University of California Davis, Davis, California 95616, USA.
- ²⁴ Center for Global Health, University of Illinois College of Medicine, Chicago, IL, USA.
- ²⁵ Subdirección de Salud Pública Gipuzkoa, San Sebastián, Spain.
- ²⁶ Instituto de Investigación Sanitaria BIODONOSTIA, San Sebastián, Spain.
- ²⁷ School of Psychology-University of the Basque Country, San Sebastián, Spain.
- ²⁸ CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain.
- ²⁹ Irish Centre for Fetal and Neonatal Translational Research, Cork University Maternity Hospital, University College Cork, Cork, Ireland.
- ³⁰ Department of Obstetrics and Gynaecology, Cork University Maternity Hospital, Cork, Ireland.
- ³¹ MRC Integrative Epidemiology Unit at the University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK.
- ³² Population Health Science, Bristol Medical School, University of Bristol, Bristol BS8 2BN, UK.
- ³³ Department of Public Health, University of Turku, Turku, Finland.
- ³⁴ Department of Environmental Immunology/Core Facility Studies, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany.
- ³⁵ Epidemiology and Environmental Health Joint Research Unit, FISABIO (Foundation for the Promotion of Health and Biomedical Research in the Valencian Region), Valencia, Spain.
- ³⁶ Division of Health Data and Digitalization, Norwegian Institute of Public Health, Oslo, Norway.
- ³⁷ Department of Epidemiology, Chair of Epidemiology and Preventive Medicine, Jagiellonian University Medical College, Krakow, Poland.
- ³⁸ Turku Centre for Biotechnology, University of Turku and Abo Akademi University, Turku, Finland.
- ³⁹ Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University, Athens, Greece.
- ⁴⁰ UCD Perinatal Research Centre, Obstetrics & Gynaecology, School of Medicine, University College Dublin, National Maternity Hospital, Dublin, Ireland.
- ⁴¹ Department of Pediatrics, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada.
- ⁴² UCD Perinatal Research Centre, School of Public Health and Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland.
- ⁴³ Department of Epidemiology, Care and Public Health Research Institute, Maastricht University, Maastricht, The Netherlands.
- ⁴⁴ National Institute of Public Health, University of Southern Denmark, Copenhagen, Denmark.
- ⁴⁵ Section of Epidemiology, Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
- ⁴⁶ Department of Rehabilitation, Nutrition and Sport; La Trobe University, Australia
- ⁴⁷ Paediatrics & Child Health, University College Cork, Cork, Ireland.
- ⁴⁸ Cork Centre for Vitamin D and Nutrition Research, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.
- ⁴⁹ Research Unit for Gynaecology and Obstetrics, Institute for Clinical Research, University of Southern Denmark, Denmark.
- ⁵⁰ Department of Population Medicine, Harvard Medical School, Harvard Pilgrim Health Care Institute, Boston, MA, USA.
- ⁵¹ Department of Public Health, Amsterdam Public Health Research Institute, Academic Medical Center, Amsterdam, The Netherlands.
- ⁵² Department of Environmental Exposures and Epidemiology, Domain of Infection Control and Environmental Health, Norwegian Institute of Public Health, Lovisenberggata 8, 0477 Oslo, Norway
- ⁵³ Department of Medical Sciences, University of Turin, Turin, Italy.
- ⁵⁴ Unit of Epidemiology, "Anna Meyer" Children's University Hospital, Florence, Italy.
- ⁵⁵ Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands.

- ⁵⁶ Novo Nordisk Foundation Center for Basic Metabolic Research, Section of Metabolic Genetics, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
- ⁵⁷ Institute of Epidemiology, Helmholtz Zentrum München-German Research Center for Environmental Health, Neuherberg, Germany.
- ⁵⁸ Norwegian Institute of Public Health, Oslo, Norway.
- ⁵⁹ Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway.
- ⁶⁰ ISGlobal, Institute for Global Health, Barcelona, Spain.
- ⁶¹ Universitat Pompeu Fabra (UPF), Barcelona, Spain.
- ⁶² Dr. von Hauner Children's Hospital, Ludwig-Maximilians-University Munich, Munich, Germany.
- ⁶³ Ib-salut, Area de Salut de Menorca, Palma, Spain.
- ⁶⁴ Department of Community Health Sciences, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada.
- ⁶⁵ Department of Environmental Medicine, Slovak Medical University, Bratislava 833 03, Slovak Republic.
- ⁶⁶ Child Health, Royal Aberdeen Children's Hospital, Aberdeen, UK.
- ⁶⁷ Research Institute, Department of Pediatrics, Marien-Hospital Wesel, Wesel, Germany.
- ⁶⁸ Bradford Institute for Health Research, Bradford Royal Infirmary, Bradford, UK.
- ⁶⁹ Department of Medical and Social Problems of Family Health, Institute of Pediatrics, Obstetrics and Gynecology, Kyiv, Ukraine.
- ⁷⁰ Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, the Netherlands.